

BIJECTIONS ON TWO VARIATIONS OF NONCROSSING PARTITIONS

JANG SOO KIM

ABSTRACT. We find bijections on 2-distant noncrossing partitions, 12312-avoiding partitions, 3-Motzkin paths, UH-free Schröder paths and Schröder paths without peaks at even height. We also give a direct bijection between 2-distant noncrossing partitions and 12312-avoiding partitions.

1. INTRODUCTION

A (set) *partition* of $[n] = \{1, 2, \dots, n\}$ is a collection of mutually disjoint nonempty subsets, called *blocks*, of $[n]$ whose union is $[n]$. We will write a partition as a sequence of blocks $(B_1, B_2, \dots, B_k)$ such that $\min(B_1) < \min(B_2) < \dots < \min(B_k)$.

There are two natural representations of a partition. Let $\pi = (B_1, B_2, \dots, B_k)$ be a partition of $[n]$. The *partition diagram* of π is the simple graph with vertex set $V = [n]$ and edge set E , where $(i, j) \in E$ if and only if i and j are in the same block which does not have an integer between them. For example, see Figure 1. The *canonical word* of π is the word $a_1 a_2 \dots a_n$, where $a_i = j$ if $i \in B_j$. For instance, the canonical word of the partition in Figure 1 is 123124412. For a word τ , a partition is called τ -*avoiding* if its canonical word does not contain a subword which is order-isomorphic to τ .

A partition is *noncrossing* if the edges of its partition diagram do not intersect. It is easy to see that a partition is noncrossing if and only if it is 1212-avoiding.

Let π be a partition and let k be a nonnegative integer. A k -*distant crossing* of π is a set of two edges (i_1, j_1) and (i_2, j_2) of the partition diagram of π satisfying $i_1 < i_2 \leq j_1 < j_2$ and $j_1 - i_2 \geq k$. A partition π is called k -*distant noncrossing* if π has no k -distant crossings. Note that 1-distant noncrossing partitions are just noncrossing partitions.

Our main objects are 2-distant noncrossing partitions and 12312-avoiding partitions. Let $\text{NC}_2(n)$ denote the set of 2-distant noncrossing partitions of $[n]$. Let $P_{12312}(n)$ denote the set of 12312-avoiding partitions of $[n]$.

Date: May 13, 2009.

2000 Mathematics Subject Classification. 05A18, 05A15.

Key words and phrases. noncrossing partitions, Motzkin paths and Schröder paths.



FIGURE 1. The partition diagram of $(\{1, 4, 8\}, \{2, 5, 9\}, \{3\}, \{6, 7\})$.

Drake and Kim [1] found the following generating function for the number of 2-distant noncrossing partitions:

$$(1) \quad \sum_{n \geq 0} \#NC_2(n)x^n = \frac{3 - 3x - \sqrt{1 - 6x + 5x^2}}{2(1 - x)}.$$

Using the kernel method, Mansour and Severini [4] found the generating function for the number of $12 \cdots k12$ -avoiding partitions of $[n]$. Interestingly, as a special case of their result, the generating function for the number of 12312-avoiding partitions of $[n]$ is the same as (1), which implies $\#NC_2(n) = \#P_{12312}(n)$. Moreover, this number also counts several kinds of lattice paths.

A *lattice path of length n* is a sequence of integer lattice points (i, j) in the xy -plane starting with $(0, 0)$ and ending with $(n, 0)$. A lattice path is *nonnegative* if it never goes below the x -axis. Since we will only consider nonnegative lattice paths, throughout this paper, we will write simply a lattice path instead of a nonnegative lattice path.

Let $L = ((x_0, y_0), (x_1, y_1), \dots, (x_k, y_k))$ be a lattice path. Then each $S_i = (x_i - x_{i-1}, y_i - y_{i-1})$ is a *step* of L . The *height* of the step S_i is defined to be y_{i-1} . Sometimes we will identify a lattice path L with the word $S_1 S_2 \dots S_k$ of its steps. Note that the number of steps is not necessarily equal to the length of the lattice path.

Let U , D and H denote an *up step*, a *down step* and a *horizontal step* respectively, i.e., $U = (1, 1)$, $D = (1, -1)$ and $H = (1, 0)$.

A *Schröder path* of length $2n$ is a lattice path of length $2n$ consisting of steps U , D and $H^2 = HH = (2, 0)$. Let $L = S_1 S_2 \dots S_k$ be a Schröder path. A *UH-pair* of L is a pair (S_i, S_{i+1}) of consecutive steps such that $S_i = U$ and $S_{i+1} = H^2$. We say that L is *UH-free* if it does not have a UH-pair. A *peak* of L is a pair (S_i, S_{i+1}) of consecutive steps such that $S_i = U$ and $S_{i+1} = D$. The *height* of a peak (S_i, S_{i+1}) is the height of $S_{i+1} = D$.

Let $SCH_{UH}(n)$ denote the set of UH-free Schröder paths of length $2n$. Let $SCH_{\text{even}}(n)$ (resp. $SCH_{\text{odd}}(n)$) denote the set of Schröder paths of length $2n$ which have no peaks of even (resp. odd) height.

A *labeled step* is a step together with an integer label. Let D_i (resp. H_i) denote a labeled down step (resp. a labeled horizontal step) with label i .

Let $CH_2(n)$ denote the set of lattice paths $L = S_1 S_2 \dots S_n$ of length n consisting of U , D_1 , D_2 , H_0 , H_1 and H_2 such that

- if $S_i = H_\ell$ or $S_i = D_\ell$, then S_i is of height at least ℓ ,
- if $S_i = H_2$ or $S_i = D_2$, then $i \geq 2$ and $S_{i-1} \in \{U, H_1, H_2\}$.

Drake and Kim [1] showed that the well known bijection φ between partitions and Charlier diagrams, see [2, 3], yields a bijection $\varphi : NC_2(n) \rightarrow CH_2(n)$. Yan [8] found a bijection $\phi : SCH_{UH}(n-1) \rightarrow P_{12312}(n)$ and a bijection between $SCH_{UH}(n)$ and $SCH_{\text{even}}(n)$.

Thus the cardinalities of $NC_2(n)$, $CH_2(n)$, $SCH_{\text{even}}(n-1)$, $SCH_{UH}(n-1)$ and $P_{12312}(n)$ are the same whose sequence appears as A007317 in [6].

Let us define the following refined sets.

- $NC'_2(n) = \{\pi \in NC_2(n) : n \text{ is not a singleton}\}$
- $CH'_2(n) = \{L \in CH_2(n) : \text{the last step of } L \text{ is } D_1\}$
- $SCH'_{\text{even}}(n) = \{L \in SCH_{\text{even}}(n) : \text{the first step of } L \text{ is } U\}$
- $SCH'_{UH}(n) = \{L \in SCH_{UH}(n) : \text{the first step of } L \text{ is } U\}$

$$\begin{array}{ccccccc}
\text{NC}'_2(n) & & & \text{SCH}'_{\text{even}}(n-1) & & & P'_{12312}(n) \\
\downarrow \varphi & & & \uparrow \iota & & & \uparrow \phi \\
\text{CH}'_2(n) & \xrightarrow{f} & \text{MOT}_3(n-2) & \xrightarrow{g} & \text{SCH}_{\text{odd}}(n-1) & \xrightarrow{h} & \text{SCH}'_{\text{UH}}(n-1)
\end{array}$$

FIGURE 2. Main bijections for $n \geq 2$.

- $P'_{12312}(n) = \{\pi \in P_{12312}(n) : 1 \text{ is not connected to } 2\}$

There are obvious bijections proving the following. For $n \geq 2$,

$$\begin{aligned}
\#\text{NC}_2(n) &= 1 + \sum_{i=2}^n \#\text{NC}'_2(i), \\
\#\text{CH}_2(n) &= 1 + \sum_{i=2}^n \#\text{CH}'_2(i), \\
\#\text{SCH}_{\text{even}}(n-1) &= 1 + \sum_{i=2}^n \#\text{SCH}'_{\text{even}}(i-1), \\
\#\text{SCH}_{\text{UH}}(n-1) &= 1 + \sum_{i=2}^n \#\text{SCH}'_{\text{UH}}(i-1), \\
\#P_{12312}(n) &= 1 + \sum_{i=2}^n \#P'_{12312}(i).
\end{aligned}$$

Thus, for $n \geq 2$, the cardinalities of the following sets are also the same, whose sequence appears as A002212 in [6] : $\text{NC}'_2(n)$, $\text{CH}'_2(n)$, $\text{SCH}'_{\text{even}}(n-1)$, $\text{SCH}'_{\text{UH}}(n-1)$ and $P'_{12312}(n)$. This number is also equal to both $\#\text{SCH}_{\text{odd}}(n-1)$ and the number of 3-Motzkin paths of length $n-2$, where a 3-Motzkin path of length n is a lattice path of length n consisting of U , D , H_0 , H_1 and H_2 .

Let $\text{MOT}_3(n)$ denote the set of 3-Motzkin paths of length n .

Our main purpose is to find bijections between these objects. For the overview of our bijections see Figure 2, where φ is the known bijection between partitions and Charlier diagrams [2, 3], and ϕ is Yan's bijection [8]. We note that our bijection g in Figure 2 is also discovered by Shapiro and Wang [5]. We also provide a direct bijection between $\text{NC}_2(n)$ and $P_{12312}(n)$.



The rest of this paper is organized as follows. In Section 2 we find bijections in Figure 2. In Section 3 we review Yan's bijection and its consequences for self containment. In Section 4 we obtain refined results following from our bijections. In Section 5 we provide a direct bijection between $\text{NC}_2(n)$ and $P_{12312}(n)$.

2. BIJECTIONS

In this section we find the bijections f, g, h and ι in Figure 2.

2.1. The bijection $f : \text{CH}'_2(n) \rightarrow \text{MOT}_3(n-2)$. Recall that $\text{CH}'_2(n)$ is the set of lattice paths $L = S_1 S_2 \cdots S_n$ of length n consisting of U, D_1, D_2, H_0, H_1 and H_2 such that

- if $S_i = H_\ell$ or $S_i = D_\ell$, then S_i is of height at least ℓ ,
- if $S_i = H_2$ or $S_i = D_2$, then $i \geq 2$ and $S_{i-1} \in \{U, H_1, H_2\}$,

FIGURE 3. An example of f_0 .FIGURE 4. Definition of f .

- $S_n = D_1$.

The second condition above is equivalent to the condition that the lattice path consists of the following combined steps for any $k \geq 0$:

$$(2) \quad UH_2^k, UH_2^k D_2, H_1 H_2^k, H_1 H_2^k D_2, H_0, D_1.$$

Let $A(n)$ denote the set of lattice paths of length n consisting of the combined steps in (2) such that H_2 does not touch the x -axis. Let $B(n)$ denote the set of 3-Motzkin paths of length n such that each H_2 touching the x -axis must occur after D , H_0 or H_2 .

We define $f_0 : A(n) \rightarrow B(n)$ as follows. Let $L \in A(n)$. Then $f_0(L)$ is defined to be the lattice path obtained from L by changing $UH_2^k D_2$ to $H_0 H_2^{k+1}$, $H_1 H_2^k D_2$ to DH_2^{k+1} and D_1 to D . It is easy to see that $f_0(L) \in B$ and f_0 is invertible. See Figure 3.

Now we define $f : \text{CH}'_2(n) \rightarrow \text{MOT}_3(n-2)$ as follows. Let $L \in \text{CH}'_2(n)$. Then L is decomposed uniquely as

$$H_0^{k_1}(UL_1 D_1) H_0^{k_2}(UL_2 D_1) \cdots H_0^{k_r}(UL_r D_1),$$

where $L_i \in A(n_i)$ for some $k_i, n_i \geq 0$ and $r \geq 1$. Then define $f(L)$ to be

$$H_2^{k_1} f_0(L_1) (H_1 H_2^{k_2+1} f_0(L_2)) (H_1 H_2^{k_3+1} f_0(L_3)) \cdots (H_1 H_2^{k_r+1} f_0(L_r)).$$

See Figure 4.

Theorem 2.1. *The map $f : \text{CH}'_2(n) \rightarrow \text{MOT}_3(n-2)$ is a bijection.*




FIGURE 5. An example of g . Odd peaks are circled. Green color is for H^2 of even height.

Proof. Each $L \in \text{MOT}_3(n-2)$ is uniquely decomposed as

$$H_2^{k_1} L_1 (H_1 H_2^{k_2+1} L_2) (H_1 H_2^{k_3+1} L_3) \cdots (H_1 H_2^{k_r+1} L_r),$$

where $L_i \in B(n_i)$ for some $k_i, n_i \geq 0$ and $r \geq 1$. Thus we have the inverse $f^{-1}(L)$ which is decomposed as

$$H_0^{k_1} (U f_0^{-1}(L_1) D_1) H_0^{k_2} (U f_0^{-1}(L_2) D_1) \cdots H_0^{k_r} (U f_0^{-1}(L_r) D_1).$$

□

2.2. The bijection $g : \text{MOT}_3(n) \rightarrow \text{SCH}_{\text{odd}}(n+1)$. We define $g : \text{MOT}_3(n) \rightarrow \text{SCH}_{\text{odd}}(n+1)$ as follows. Let $L \in \text{MOT}_3(n)$. Then $g(L)$ is the lattice path obtained from L by doing the following.

- (1) Change U to UU , D to DD , H_0 to H^2 , H_1 to DU , and H_2 to UD .
- (2) Add U at the beginning and D at the end.
- (3) Change all the consecutive steps UD which form a peak of odd height to H^2 .

See Figure 5 for an example of g .

Theorem 2.2. *The map $g : \text{MOT}_3(n) \rightarrow \text{SCH}_{\text{odd}}(n+1)$ is a bijection.*

Proof. Clearly the first and the second steps in the construction of g are invertible. The third step is also invertible because every step H^2 of even height always comes from a peak of odd height. Thus g is invertible. □

2.3. The bijection $h : \text{SCH}_{\text{odd}}(n) \rightarrow \text{SCH}'_{\text{UH}}(n)$. Let $L = S_1 S_2 \cdots S_k$ be a Schröder path. Let $S_i = U$ be an up step. Then there is a unique down step $S_j = D$ such that $i < j$ and $S_{i+1} S_{i+2} \cdots S_{j-1}$ is a (possibly empty) lattice path. We call such S_j the down step corresponding to S_i . We also call S_i the up step corresponding to S_j .




FIGURE 6. The essence of h_0 . Green (resp. Red) color is for UH-pairs with H^2 of even (resp. odd) height. Odd peaks are circled. The lattice path L' is not empty.

Let $L = S_1 S_2 \cdots S_k \in \text{SCH}_{\text{odd}}(n)$. Recall that a UH-pair of L is a pair (S_i, S_{i+1}) such that $S_i = U$ and $S_{i+1} = H^2$. Let (S_i, S_{i+1}) be a UH-pair and let S_j be the down step corresponding to S_i . Then we define the function ξ as follows.

$$\xi(S_i, S_{i+1}) = \begin{cases} i, & \text{if } S_{i+1} \text{ is of even height;} \\ j, & \text{if } S_{i+1} \text{ is of odd height.} \end{cases}$$

If L is not UH-free, we define the ξ -maximal UH-pair of L to be the UH-pair (S_i, S_{i+1}) satisfying the following: for any UH-pair $(S_{i'}, S_{i'+1})$ with $i' \neq i$, we have $\xi(S_i, S_{i+1}) > \xi(S_{i'}, S_{i'+1})$.

Let $L \in \text{SCH}_{\text{odd}}(n)$. Assume that L is not UH-free. Let (S_i, S_{i+1}) be the ξ -maximal UH-pair of L . Let S_j be the down step corresponding to S_i . Then we define $h_0(L)$ as follows.

- (1) If S_{i+1} is of even height, then $h_0(L)$ is the lattice path obtained from L by replacing $S_i S_{i+1}$ with UUH .
- (2) If S_{i+1} is of odd height, then let $L' = S_{i+2} S_{i+3} \cdots S_{j-1}$.
 - (a) If L' is empty, i.e., $j = i + 2$, then $h_0(L)$ is the lattice path obtained from L by replacing $S_i S_{i+1} S_{i+2}$ with $H^2 U D$.
 - (b) If L' is not empty, then $h_0(L)$ is the lattice path obtained from L by replacing $S_i S_{i+1} \cdots S_j$ with $U L' D U D$.

See Figure 6.

Now we define $h : \text{SCH}_{\text{odd}}(n) \rightarrow \text{SCH}'_{\text{UH}}(n)$ as follows. Let $L \in \text{SCH}_{\text{odd}}(n)$ and $L_0 = L$. Then we define $L_i = h_0(L_{i-1})$ for $i \geq 1$ if L_{i-1} is not UH-free. Since the number of UH-free pairs of L_i is one less than that of L_{i-1} , or they are the same and

$$\xi(\text{the maximal UH-pair of } L_i) < \xi(\text{the maximal UH-pair of } L_{i-1}),$$

we always get L_r which is UH-free for some r . We define $h(L)$ to be L_r if L_r does not start with H^2 ; and the lattice path obtained from L_r by replacing H^2 with UD otherwise. For example, see Figure 7.

Theorem 2.3. *The map $h : \text{SCH}_{\text{odd}}(n) \rightarrow \text{SCH}'_{\text{UH}}(n)$ is a bijection.*

Proof. In the procedure of h , the odd peaks are constructed from right to left. Since h_0 is invertible, so is h . $\square$





FIGURE 7. An example of h . Green (resp. Red) color is for UH-pairs with H^2 of even (resp. odd) height. Odd peaks are circled. Dashed arrows indicate the down steps corresponding to the up steps.

2.4. **The bijection $\iota : \text{SCH}_{\text{odd}}(n) \rightarrow \text{SCH}'_{\text{even}}(n)$.** Let $L = S_1 S_2 \cdots S_k \in \text{SCH}_{\text{odd}}(n)$. Then $\iota(L)$ is defined as follows.

- (1) If $S_k = H^2$, then $\iota(L) = U S_1 \cdots S_{k-1} D$.
- (2) If $S_k = D$, then let S_i be the up step corresponding to S_k and we define $\iota(L) = U S_1 \cdots S_{i-1} D S_{i+1} \cdots S_{k-1}$.

FIGURE 8. The map ι .

See Figure 8.

Then $\iota(L) \in \text{SCH}'_{\text{even}}(n)$. Clearly, $\iota : \text{SCH}_{\text{odd}}(n) \rightarrow \text{SCH}'_{\text{even}}(n)$ is a bijection.

3. YAN'S BIJECTION

In this section we describe Yan's bijection $\phi : \text{SCH}_{\text{UH}}(n-1) \rightarrow P_{12312}(n)$.

Let $L \in \text{SCH}_{\text{UH}}(n-1)$. Then we label each step of L as follows.

- (1) Find all the peaks of L , and label the up step of the i -th leftmost peak with $i+1$ for $i = 1, 2, \dots$
- (2) We label all the unlabeled up steps and the horizontal steps with the largest integers which are left to the steps. If there is no integer which is left to such a step, then we label it with 1.
- (3) For each down step D , we label it with the same integer which is the label of the up step corresponding to D .

Then ϕ is defined to be the word obtained by reading the labels of down steps and horizontal steps of L from left to right and by adding 1 at the beginning. See Figure 9 for an example of this map.

Let $\pi = (B_1, B_2, \dots, B_r)$ be a partition. A pair (B_i, a) of a block and an integer is called *inversion* if $a > \min(B_i)$ and $a \in B_j$ and for some $j < i$. Let $\text{inv}(B_i)$ denote the number of inversions of form (B_i, a) .

From the definition of ϕ , we get the following immediately for L and π with $\phi(L) = \pi$.

- The first step of L is a horizontal step if and only if 1 is connected to 2 in π
- If $\pi = (B_1, \dots, B_k)$, then L has $k-1$ peaks and their heights are $\text{inv}(B_2) + 1, \text{inv}(B_3) + 1, \dots, \text{inv}(B_k) + 1$.

Thus ϕ is indeed a bijection between $\text{SCH}'_{\text{UH}}(n-1)$ and $P'_{12312}(n)$.

Using this bijection Yan proved the following theorem.

Theorem 3.1. [8] *Let $n, k \geq 1$. Then the number of 12312-avoiding partitions of $[n+1]$ with $k+1$ blocks is equal to*

$$\sum_{i=k}^n \frac{1}{i} \binom{i}{k-1} \binom{i}{k} \binom{n}{i}.$$




FIGURE 9. Yan's bijection



FIGURE 10. All the five crossings are high crossings.

4. REFINED RESULTS

Let $\pi \in \text{NC}'_2(n)$. Let $\{(a, i+1), (i, b)\}$ be a crossing of π . Let A and B be the blocks containing a and b respectively. Let $r = \min(\max(A), \max(B))$. This crossing is called *low crossing* if it satisfies the following:

- $(a, i+1)$ is not nested by another edge,
- $i, i+1, \dots, r$ are contained in B and A alternatively, that is, $i \in B, i+1 \in A, i+2 \in B$, and so on.

Otherwise, it is called *high crossing*. For example, see Figure 10 and Figure 11.

Let $L \in \text{CH}'_2(n)$. A step D_2 of L is *low* if its height is 2. Otherwise it is *high*. A step H_2 of L is *low* if the first step right to it, which is not H_2 , is a low D_2 . Otherwise it is *high*. A *high step* is either a high D_2 or a high H_2 .




FIGURE 11. All the five crossings are low crossings.

Let us define the following refined sets.

- $\text{NC}'_2(n, k) = \{\pi \in \text{NC}'_2(n) : \pi \text{ has } k \text{ high crossings}\}$
- $\text{CH}'_2(n, k) = \{L \in \text{CH}'_2(n) : L \text{ has } k \text{ high steps}\}$
- $\text{MOT}_3(n-2, k) = \{L \in \text{MOT}_3(n-2) : L \text{ has } k \text{ } H_2\text{'s of height } > 0\}$
- $\text{SCH}_{\text{odd}}(n-1, k) = \{L \in \text{SCH}_{\text{odd}}(n-1) : L \text{ has } k \text{ peaks of even height } > 2\}$
- $\text{SCH}'_{\text{UH}}(n-1, k) = \{L \in \text{SCH}'_{\text{UH}}(n-1) : L \text{ has } k \text{ peaks of even height } > 2\}$
- $P'_{12312}(n, k) = \{\pi \in P'_{12312}(n) : \pi \text{ has } k \text{ blocks } B_i \text{ with odd } \text{inv}(B_i) > 1\}$

It is easy to check that our maps in Figure 2 induce bijections on the sets described above.

Let $p(n, k)$ be the number of one of the objects described above. Let $a(n, k)$ be the number of 3-Motzkin paths of length n with k H_2 steps. Let $b(n, k)$ be the number of 3-Motzkin paths of length n with k H_2 steps of height at least 1. Let

$$\begin{aligned} A(x, y) &= \sum_{n \geq 0} \sum_{m \geq 0} a(n, m) x^n y^m, \\ B(x, y) &= \sum_{n \geq 0} \sum_{m \geq 0} b(n, m) x^n y^m, \\ F(x, y) &= \sum_{n \geq 2} \sum_{m \geq 0} p(n, m) x^n y^m. \end{aligned}$$

Then it is easy to see that

$$\begin{aligned} A(x, y) &= 1 + (2 + y)x A(x, y) + x^2 A(x, y)^2, \\ B(x, y) &= 1 + 3xB(x, y) + x^2 B(x, y)A(x, y), \\ F(x, y) &= x^2 \cdot B(x, y). \end{aligned}$$

Solving these equations, we get

$$F(x, y) = \frac{2x^2}{1 - (4 - y)x + \sqrt{1 - (4 + 2y)x + (4y + y^2)x^2}}.$$

If $y = 0$, then

$$\begin{aligned} F(x, 0) &= \frac{2x^2}{1 - 4x + \sqrt{1 - 4x}} = \frac{2x^2}{\sqrt{1 - 4x}} \left(\frac{1}{1 + \sqrt{1 - 4x}} \right) \\ &= \frac{2x^2}{\sqrt{1 - 4x}} \left(\frac{1 - \sqrt{1 - 4x}}{4x} \right) = \frac{x}{2} \left((1 - 4x)^{-\frac{1}{2}} - 1 \right). \end{aligned}$$

Note that

$$(1 - 4x)^{-\frac{1}{2}} = \sum_{n \geq 0} \binom{-\frac{1}{2}}{n} (-4x)^n = \sum_{n \geq 0} \binom{2n}{n} x^n.$$

Thus we have the following.

Proposition 4.1. *If $n \geq 2$, then*

$$p(n, 0) = \frac{1}{2} \binom{2n-2}{n-1}.$$

In fact, we can prove Proposition 4.1 bijectively as follows. Recall that $p(n, k)$ is equal to the number of elements in

$$\text{SCH}_{\text{odd}}(n-1, k) = \{L \in \text{SCH}_{\text{odd}}(n-1) : L \text{ has } k \text{ peaks of even height } > 2\}.$$

Thus it is sufficient to show the following:

$$\frac{1}{2} \binom{2n}{n} = \#\text{SCH}_{\text{odd}}(n, 0).$$

If $L \in \text{SCH}_{\text{odd}}(n, 0)$, then the height of each peak of L is 2. Let L' be the Schröder path obtained from L by changing each horizontal step H^2 whose height is not 1 to a peak UD . Then L' is a Schröder path such that each horizontal step is of height 1. This gives us a bijection between $\text{SCH}_{\text{odd}}(n, 0)$ and $\text{SCH}_1(n)$, the set of all Schröder paths of length $2n$ such that the height of each horizontal step is 1.

The following proposition finishes our bijective proof of Proposition 4.1.

Proposition 4.2. *There is a bijection between $\text{SCH}_1(n)$ and the set of paths from $(0, 0)$ to (n, n) with steps $(1, 0)$ and $(0, 1)$ starting with a step $(1, 0)$.*

Proof. We can consider $L \in \text{SCH}_1(n)$ as a path from $(0, 0)$ to (n, n) with steps $(1, 0)$, $(0, 1)$ and $(1, 1)$ that never goes below the line $y = x$. Then L has steps $(1, 1)$ only on the line $y = x + 1$. Let $L = (S_1, S_2, \dots, S_m)$ and let $S_{i_1}, S_{i_2}, \dots, S_{i_\ell}$ be the steps $(1, 1)$. Then we change S_{i_ℓ} to two consecutive steps $S = (1, 0)$, $S' = (0, 1)$, and then reflect the path starting from S' to the last step along the line $y = x$. We do the same thing with $S_{i_{\ell-1}}$ on the resulting path, and so on. Then at the end, we get a path with steps $(1, 0)$ and $(0, 1)$ starting with $(1, 0)$. It is easy to see that this procedure is reversible. $\square$

By finding the generating function, Yan [7] showed that the number of UH-free Schröder paths of length n without horizontal steps of height greater than 1 is equal to

$$(3) \quad \frac{1}{2} \left(1 + \sum_{i=0}^n \binom{2i}{i} \right).$$

The following proposition gives us a bijective proof of (3).

Proposition 4.3. *There is a bijection between $\text{SCH}_1(n)$ and the set of UH-free Schröder paths of length n , starting with an up step, and with no horizontal steps of height greater than 1.*

Proof. Let L be a UH-free Schröder path with the conditions. Then we can decompose L as follows:

$$L = (UL_1DH^{k_1})(UL_2DH^{k_2}) \cdots (UL_\ell DH^{k_\ell}),$$

where $k_i \geq 0$ and L_i is a (possibly empty) Schröder path with no horizontal steps of height greater than 0 which does not start with a horizontal step. Let

$$L' = (UH^{k_1}L_1D)(UH^{k_2}L_2D) \cdots (UH^{k_\ell}L_\ell D).$$

Then $L' \in \text{SCH}_1(n)$. Clearly, this is invertible. $\square$

5. A DIRECT BIJECTION BETWEEN $\text{NC}_2(n)$ AND $P_{12312}(n)$

Now we have a bijection $\phi \circ h \circ g \circ f \circ \varphi : \text{NC}_2(n) \rightarrow P_{12312}(n)$, see Figure 2. Since both $\text{NC}_2(n)$ and $P_{12312}(n)$ are partitions with some conditions, it is natural to ask a direct bijection between them. In this section we find such a direct bijection.

From now on, we will identify a partition in $P_{12312}(n)$ with its canonical word.

A *marked* partition is a partition with marking some integers. Each marked integer is called *active*. Similarly a *marked* word is a word with marking some letters. Each marked letter is called *active*.

Let $\pi \in \text{NC}_2(n)$. For $i \in [n]$, let T_i be the marked partition of $[i]$ obtained from π by removing all the integers greater than i and by marking integers which are connected to an integer greater than i .

Using the sequence $\emptyset = T_0, T_1, T_2, \dots, T_n = \pi$ of marked partitions, we define a sequence of marked words $\mathbf{w}_0, \mathbf{w}_1, \mathbf{w}_2, \dots, \mathbf{w}_n$ as follows.

Let $\mathbf{w}_0$ be the empty word.

For $1 \leq i \leq n$, $\mathbf{w}_i$ is defined as follows.

- (1) If i is not connected to any integer in T_i , then $\mathbf{w}_i = \mathbf{w}_{i-1}m$, where $m = \max(\mathbf{w}_{i-1}) + 1$. Otherwise, i is connected to either the largest active integer or the second largest active integer of T_{i-1} .
 - If i is connected to the largest active integer of T_{i-1} , then let $\mathbf{w}_i = \mathbf{w}_{i-1}a_1$, where a_1 is the rightmost active letter of $\mathbf{w}_{i-1}$. And then we make the active letter a_1 inactive.
 - If i is connected to the second largest active integer of T_{i-1} , then let $\mathbf{w}_i = \mathbf{w}_{i-1}a_2$, where a_2 is the second rightmost active letter of $\mathbf{w}_{i-1}$. The second rightmost active letter of $\mathbf{w}_{i-1}$ remains active, however, we make the rightmost active letter of $\mathbf{w}_{i-1}$ inactive in $\mathbf{w}_i$.
- (2) If i is active in T_i , then we find the largest letters in $\mathbf{w}_i$ and make the leftmost letter among them active.

For example, see Figure 12. It is easy to check that $\mathbf{w}_n$ is 12312-avoiding.

When we know $\mathbf{w}_n$, we can reverse this procedure. First, we find $\mathbf{w}_i$'s as follows.

For $1 \leq i \leq n$, $\mathbf{w}_{i-1}$ is obtained from $\mathbf{w}_i$ as follows.

- (1) Let $m = \max(\mathbf{w}_i)$. Let t be the last letter of $\mathbf{w}_i$.
- (2) If the leftmost m is active in $\mathbf{w}_i$, then make it inactive.
- (3) If t appears only once in $\mathbf{w}_i$ (equivalently t is greater than any other letters in $\mathbf{w}_i$), then we simply remove t . Otherwise, find the leftmost t in $\mathbf{w}_i$.
 - If the leftmost t is inactive, then we remove the last letter t and make the leftmost t active.
 - If the leftmost t is active, then we must have $t < m$ since we have made the leftmost m inactive. In this case we remove the last t , and make the leftmost t still active and the leftmost m active.

Now we construct $T_0, T_1, \dots, T_n$ as follows.

Let $T_0 = \emptyset$. For $1 \leq i \leq n$, T_i is obtained as follows.

- (1) First, let T_i be the marked partition obtained from T_{i-1} by adding i .
- (2) Let t be the last letter of $\mathbf{w}_i$.
- (3) If t is equal to the rightmost (resp. the second rightmost) active letter of $\mathbf{w}_{i-1}$, then connect i to the largest (resp. the second largest) active integer, say j , of T_{i-1} , and make j inactive.




FIGURE 12. T_i 's and corresponding w_i 's. Active integers and active letters are circled.

- (4) Let $m = \max(w_i)$. If the leftmost m is active in w_i , then make i active in T_i .

It is easy to check that this is the inverse map. Thus we get the following theorem.

Theorem 5.1. *For $\pi \in \text{NC}_2(n)$, the map $\pi \mapsto w_n$ is a bijection from $\text{NC}_2(n)$ to $P_{12312}(n)$.*

Since this bijection preserves the number of blocks, by Theorem 3.1, we get the following.

Corollary 5.2. *Let $n, k \geq 1$. Then the number of 2-distant noncrossing partitions of $[n+1]$ with $k+1$ blocks is equal to*

$$\sum_{i=k}^n \frac{1}{i} \binom{i}{k-1} \binom{i}{k} \binom{n}{i}.$$

REFERENCES

- [1] Dan Drake and Jang Soo Kim. k -distant crossings and nestings of matchings and partitions. <http://arxiv.org/abs/0812.2725>

- [2] P. Flajolet. Combinatorial aspects of continued fractions. *Discrete Math.*, 32(2):125-161, 1980.
- [3] Anisse Kasraoui and Jiang Zeng. Distribution of crossings, nestings and alignments of two edges in matchings and partitions. *Electron. J. Combin.*, 13(1):Research Paper 33, 12 pp. (electronic), 2006.
- [4] T. Mansour and S. Severini. Enumeration of $(k, 2)$ -noncrossing partitions. *Discrete Math.*, 308(20):4570-4577, 2008.
- [5] Louis W. Shapiro and Carol J. Wang. A bijection between 3-Motzkin paths and Schröder paths with no peaks at odd height. *Journal of Integer Sequences*, 12:Article 09.3.2, 2009
- [6] N. J. A. Sloane. The on-line encyclopedia of integer sequences.
<http://www.research.att.com/~njas/sequences/>
- [7] Sherry H.F. Yan. Schröder paths and pattern avoiding partitions.
<http://arxiv.org/abs/0805.2465v1>
- [8] Sherry H.F. Yan. Schröder paths and pattern avoiding partitions.
<http://arxiv.org/abs/0805.2465>

E-mail address: jskim@lri.fr